

Ising transition in dimerized XY quantum spin chain

Fei Ye Guo-Hui Ding Bo-Wei Xu

Department of Physics, Shanghai Jiao Tong University, Shanghai 200030, China

(February 1, 2008)

Abstract

In the present paper, we proposed a simple spin-1/2 model which provides a exactly solvable example to study the Ising criticality with central charge $c = 1/2$. By mapping it onto the real Majorana fermions, the Ising critical behavior is explored explicitly, although its bosonized form is not the double frequency sine-Gordon model.

PACS: 75.10.Jm

Keywords: Quantum spin; Ising transition

Recently Delfino and Mussardo have shown that in the double frequency sine-Gordon (DSG) model there exists a quantum critical line where the DSG model displays an Ising criticality with central charge $c=1/2$, which has stimulated theoretical interest [1–4]. This phenomenon is quite general in the various spin chains and other quantum one-dimensional (1D) systems. By constructing the mappings between the DSG model and the deformed Ashkin-Teller (DAT) model, Fabrizio *et al* investigated the Ising transition in details [2]. They gave the relation between the original DSG operators and those from the DAT model and identified the degrees of freedom in the DAT model that become critical. They also demonstrated the efficiency of their approach to describe Ising transition in some physical realizations of the DSG model, such as the Heisenberg chain in a staggered magnetic field and the 1D Hubbard model with a staggered potential and accomplished a description of the critical properties of the Ising transition. In our previous study [4], we analysed the DSG model with the renormalisation group(RG) technique. Through the one loop β equation of the DSG Hamiltonian, two Ising fixed points and the topology of the RG flows are obtained. Along the critical line, the DSG model flows from the central charge $c=1$ Gaussian fixed point to the $c=1/2$ Ising fixed point according to the Zamolodchikov's c -theorem [5,6].

In this paper we will show the mentioned phenomenon also occurs in a simpler dimerized XY (DXY) model. Such a model provides us an exactly solvable example to show how the Ising criticality with central charge $c = 1/2$ occurs, since its low energy behavior can be described by two real Majorana fermions, one of which becomes massless on some critical line.

The Hamiltonian of the DXY model reads

$$H = \sum_n [(1 + \gamma)S_n^x S_{n+1}^x + (1 - \gamma)S_n^y S_{n+1}^y + \delta(-1)^n (S_n^x S_{n+1}^x + S_n^y S_{n+1}^y)] , \quad (1)$$

where S_n are the spin operators on the lattice sites n , γ is a parameter describing the anisotropic XY model, and δ being the coupling constant of the dimerized interaction.

It is easy to show that both the anisotropy term and the dimerization term in the DXY model will open a gap 2γ and 2δ , if acting alone, respectively. However if they coexist, as

we shall show, the excitation gap will vanish by fine tuning of the parameters δ and γ in the DXY model. This implies the Ising criticality with central charge $c = 1/2$, which is revealed obviously by transforming the spin operators in DXY model into the real Majorana fields.

Under Jordan-Wigner transformation

$$\begin{aligned} S_n^+ &= c_n^\dagger \exp(i\pi \sum_{l < n} c_l^\dagger c_l) \\ S_n^- &= c_n \exp(-i\pi \sum_{l < n} c_l^\dagger c_l) \\ S_n^z &= c_n^\dagger c_n - \frac{1}{2}, \end{aligned} \quad (2)$$

which maps spins onto spinless fermions, Eq.(1) becomes

$$\begin{aligned} H &= \frac{1}{2} \sum_{n=1}^{N-1} [(1 + (-1)^n \delta)(c_n^\dagger c_{n+1} + h.c.) + \gamma(c_n^\dagger c_{n+1}^\dagger + h.c.)] \\ &\quad - \frac{1}{2} [c_N^\dagger c_1 + \gamma c_N^\dagger c_1^\dagger + \delta c_N^\dagger c_1 + h.c.] \exp(i\pi \sum_{l=1}^N c_l^\dagger c_l), \end{aligned} \quad (3)$$

where c_n and $c_n^\dagger$ denote the annihilation and creation operators for a fermion on site n , respectively. In Eq.(4), $\exp(i\pi \sum_{l=1}^N c_l^\dagger c_l)$ is conserved and has eigenvalues ± 1 which correspond to the cyclic condition and the anticyclic condition, i.e., $c_1 = c_{N+1}$ and $c_1 = -c_{N+1}$, respectively. In the following, we shall focus on the cyclic condition.

For small δ and γ , one can express the fermi operators in terms of the right-moving(R) and left-moving(L) operators near the fermi points

$$c_n = (-i)^n R_n + i^n L_n, \quad (4)$$

the Hamiltonian density of Eq.(4) then can be represented in the continuum limit ($x = na$) as

$$\begin{aligned} \mathcal{H} &= -iR^\dagger(x)\partial_x R(x) + iL^\dagger(x)\partial_x L(x) \\ &\quad - i\gamma[R^\dagger(x)L^\dagger(x) - L(x)R(x)] + i\delta[R^\dagger(x)L(x) - L^\dagger(x)R(x)]. \end{aligned} \quad (5)$$

For our purposes, we introduce two real Majorana fermi fields $\xi_{R(L)}^{(i)}$ ($i = 1, 2$)

$$R = \frac{1}{\sqrt{2}}(\xi_R^{(1)} + i\xi_R^{(2)}) \quad L = \frac{1}{\sqrt{2}}(\xi_L^{(1)} + i\xi_L^{(2)}) \quad (6)$$

and rewrite Eq.(6) as

$$\mathcal{H} = -\frac{i}{2} \sum_{a=1}^2 (\xi_R^{(a)} \partial_x \xi_R^{(a)} - \xi_L^{(a)} \partial_x \xi_L^{(a)}) - i(\gamma - \delta) \xi_R^{(1)} \xi_L^{(1)} + i(\gamma + \delta) \xi_R^{(2)} \xi_L^{(2)} . \quad (7)$$

Eq.(7) is the standard form of the Hamiltonian density of two different Majorana fields with the masses $m_1 = \gamma - \delta$ and $m_2 = -\gamma - \delta$, respectively. The Ising criticality is reached when $\gamma = \pm\delta$, one of the masses vanishes. At this point, the low energy behavior is dominated by the massless Majorana fermion, which is Ising like, although the bosonized form of the DXY model is not the DSG model.

Since the Hamiltonian of Eq.(4) has a simple quadratic form of fermi operators, it can be diagonalized exactly. From the exact excitation spectrum, one can see that the energy gap vanishes on the critical line $\delta = \pm\gamma$, which confirms the above perturbation theory and indicates the Ising criticality.

In order to diagonalize the Hamiltonian of Eq.(4), we first consider a more general quadratic form in fermi operators

$$H = \sum_{i,j} c_i^\dagger A_{ij} c_j + \frac{1}{2} (c_i^\dagger B_{ij} c_j^\dagger + h.c.) . \quad (8)$$

The Hermiticity of H requires that $\mathbf{A}$ be a Hermitian matrix, while the anticommutation relationship among the fermi operators require that $\mathbf{B}$ be an antisymmetric matrix. Both $\mathbf{A}$ and $\mathbf{B}$ are assumed to be real.

Such a Hamiltonian can be reduced to the diagonal form [7]

$$H = \sum_k E_k d_k^\dagger d_k + \frac{1}{2} (Tr A - \sum_k E_k) , \quad (9)$$

by a linear transformation

$$d_k = \sum_i \frac{\phi_{ki} + \psi_{ki}}{2} c_i + \frac{\phi_{ki} - \psi_{ki}}{2} c_i^\dagger , \quad (10)$$

$$d_k^\dagger = \sum_i \frac{\phi_{ki} + \psi_{ki}}{2} c_i^\dagger + \frac{\phi_{ki} - \psi_{ki}}{2} c_i , \quad (11)$$

where ϕ_{ki} and ψ_{ki} , considered as real, are to be determined. This transformation should ensure that the fermi operators $d_k^\dagger$ and d_k obey the anticommutation rules $\{d_k, d_l^\dagger\} = \delta_{kl}$ and $\{d_k, d_l\} = 0$. By substituting Eqs.(10,11) into

$$[d_k, H] - E_k d_k = 0 , \quad (12)$$

one can easily find

$$\phi_k(\mathbf{A} - \mathbf{B}) = E_k \psi_k \quad (13)$$

$$\psi_k(\mathbf{A} + \mathbf{B}) = E_k \phi_k , \quad (14)$$

in an matrix notation. Then it is straightforward from Eqs.(13,14) to see that the N-component vectors ϕ_k and ψ_k satisfy the following two eigenfunctions

$$\phi_k(\mathbf{A} - \mathbf{B})(\mathbf{A} + \mathbf{B}) = E_k^2 \phi_k \quad (15)$$

and

$$\psi_k(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B}) = E_k^2 \psi_k , \quad (16)$$

respectively, belonging to the eigenvalue E_k^2 . For cyclic matrices $\mathbf{A}$ and $\mathbf{B}$, it is easy to solve the eigenfunctions (15) or (16). In our case, the relevant matrices are

$$\mathbf{A} = \frac{1}{2} \begin{pmatrix} 0 & 1-\delta & 0 & 0 & \cdots & 0 & 1+\delta \\ 1-\delta & 0 & 1+\delta & 0 & \cdots & 0 & 0 \\ 0 & 1+\delta & 0 & 1-\delta & \cdots & 0 & 0 \\ 0 & 0 & 1-\delta & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1-\delta \\ 1+\delta & 0 & 0 & 0 & \cdots & 1-\delta & 0 \end{pmatrix} \quad (17)$$

and

$$\mathbf{B} = \frac{1}{2} \begin{pmatrix} 0 & \gamma & 0 & 0 & \cdots & 0 & -\gamma \\ -\gamma & 0 & \gamma & 0 & \cdots & 0 & 0 \\ 0 & -\gamma & 0 & \gamma & \cdots & 0 & 0 \\ 0 & 0 & -\gamma & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 & \gamma \\ \gamma & 0 & 0 & 0 & \cdots & -\gamma & 0 \end{pmatrix} . \quad (18)$$

Then, the Hermitian matrix $(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B})$ reads

$$\frac{1}{2} \begin{pmatrix} 1 + (\delta - \gamma)^2 & 0 & \frac{1}{2}[1 - (\delta - \gamma)^2] & 0 & \cdots & \frac{1}{2}[1 - (\delta - \gamma)^2] & 0 \\ 0 & 1 + (\delta + \gamma)^2 & 0 & \frac{1}{2}[1 - (\delta + \gamma)^2] & \cdots & 0 & \frac{1}{2}[1 - (\delta + \gamma)^2] \\ \frac{1}{2}[1 - (\delta - \gamma)^2] & 0 & 1 + (\delta - \gamma)^2 & 0 & \cdots & 0 & 0 \\ 0 & \frac{1}{2}[1 - (\delta + \gamma)^2] & 0 & 1 + (\delta + \gamma)^2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{1}{2}[1 - (\delta - \gamma)^2] & 0 & 0 & 0 & \cdots & 1 - (\delta - \gamma)^2 & 0 \\ 0 & \frac{1}{2}[1 - (\delta + \gamma)^2] & 0 & 0 & \cdots & 0 & 1 - (\delta + \gamma)^2 \end{pmatrix}, \quad (19)$$

which are cyclic with period 2. To show how to calculate the eigenvalue of the matrix of Eq.(19), we consider a general cyclic matrix with period 2, which has the form

$$\mathbf{M} = \begin{pmatrix} a_1 & a_2 & a_3 & a_4 & \cdots & a_N \\ b_1 & b_2 & b_3 & b_4 & \cdots & b_N \\ a_{N-1} & a_N & a_1 & a_2 & \cdots & a_{N-2} \\ b_{N-1} & b_N & b_1 & b_2 & \cdots & b_{N-2} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ b_3 & b_4 & b_5 & b_6 & \cdots & b_2 \end{pmatrix}. \quad (20)$$

It has eigenvectors

$$V_k = (\epsilon, \xi\epsilon^2, \epsilon^3, \xi\epsilon^4, \dots, \xi\epsilon^N)^T \quad (21)$$

with $\epsilon = \exp(ik)$, $k = (2n\pi)/N$ ($n = 0, 1, 2, \dots, N$) and parameter ξ to be determined, corresponding the eigenvalue λ_k . The real ϕ_k and ψ_k can be extracted from the real or imaginary part of V_k , since the matrice $(\mathbf{A} \pm \mathbf{B})(\mathbf{A} \mp \mathbf{B})$ are real in our case. Substituting Eq.(21) into the eigenfunction $(\mathbf{M} - \lambda_k \mathbf{I})V_k = 0$, one obtains

$$\begin{aligned} \lambda_k &= a_1 + \xi a_2 \epsilon + \cdots + \xi a_N \epsilon^N \\ &= (\xi \epsilon)^{-1} (b_1 + b_2 \xi \epsilon + b_3 \epsilon^2 + \cdots + b_N \xi \epsilon^{N-1}). \end{aligned} \quad (22)$$

For the matrix $(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B})$ we are interested in, the eigenvalues E_k^2 are $\cos^2 k + (\delta + \gamma)^2 \sin^2 k$ and $\cos^2 k + (\delta - \gamma)^2 \sin^2 k$.

Consequently, the energy spectrum of the Hamiltonian of Eq.(4) are written as

$$E_k^{(+)} = \pm \sqrt{\cos^2 k + (\delta + \gamma)^2 \sin^2 k} , \quad (23)$$

$$E_k^{(-)} = \pm \sqrt{\cos^2 k + (\delta - \gamma)^2 \sin^2 k} . \quad (24)$$

Eqs.(23,24) indicate that the energy spectrum has two branches, one with a gap $2|\delta + \gamma|$ and the other with a gap $2|\delta - \gamma|$, which correspond to the two massive Majorana fermions, respectively. The exact solution shows unambiguously that one of the excitation gap vanishes when $\delta = \pm\gamma$, where the Ising phase transition occurs.

In summary, in the present paper, we propose a spin-1/2 model DXY chain which provides a simple example to study the Ising criticality with central charge $c = 1/2$. By mapping it onto the real Majorana fermions and diagonalizing it exactly, the Ising critical behavior is explored explicitly, although its bosonized form is not the DSG model. It seems that the Ising phase transition is quite a general feature for a variety of spin chain systems besides those which can be transformed into the DSG model.

The project is supported by the National Natural Science Foundation of China(Grant No.19975031) and the RFDP(Grant No.199024833).

REFERENCES

- [1] G.Delfino and G.Mussardo *Nucl. Phys. B* **516** (1998) 675
- [2] M.Fabrizio, A.O.Gogolin and A.A.Nersesyan *Nucl. Phys. B* **580** (2000) 647
- [3] Z.Bajnok, L.Palla, G.Takacs and F.Wagner *Nucl. Phys. B* **601** (2001) 503
- [4] F.Ye, G.H.Ding and B.W.Xu cond-mat/0104321
- [5] A.B.Zamolodchikov *JETP. Lett.* **43** (1986) 730
- [6] J.L.Cardy 1988 *Lectures delivered at Les Houches Summer School on Fields, strings and Statistical Mechanics*
- [7] E.Lieb, T.Schultz and D.Mattis, *Ann.Phys.* **16** (1961) 407